

Exercises for Graham's lectures

1. The primary ideals of \mathbb{Z} are precisely the ideals (p^m) , where p is a prime integer and $m \geq 1$. The radical of such an ideal is just (p) .
2. For a local ring (R, \mathfrak{m}, k) and a module M , elements $a_1, \dots, a_n \in M$ form a minimal generating set for M if and only if their images in $M/\mathfrak{m}M$ form a vector space basis.
3. Suppose x is R -regular. Prove that $\text{Tor}_1^R(M, R/xR)$ is isomorphic to the submodule of M killed by x .
4. Show that the previous exercise is false if x is not assumed R -regular.
5. Show that a finitely generated module M over a local ring (R, \mathfrak{m}) has finite length (i.e., a composition series) if and only if $\text{Supp}_R(M) = \{\mathfrak{m}\}$.
6. Let (R, \mathfrak{m}) be local and M, N two nonzero finitely generated modules such that $\text{pd}_R M = n < \infty$. Show that $\text{Ext}_R^n(M, N) \neq 0$.
7. For $n \geq 2$, let $R = k[x_1, \dots, x_n]$ and $I = (x_1x_2, x_2x_3, \dots, x_nx_1)$ (the edge ideal of the n -cycle). Find all n such that R/I is Cohen-Macaulay.
8. Let (R, \mathfrak{m}) be local and M, N finitely generated. Suppose that $\text{Tor}_i^R(M, N) = 0$ for all $i > 0$. Prove that $\text{pd}_R(M \otimes_R N) = \text{pd}_R M + \text{pd}_R N$.
9. Give an example to show that the previous exercise is false without the vanishing assumption on Tor . In fact, $M \otimes_R N$ might not even have finite projective dimension.
10. Prove that the tensor product of two complexes is a complex.
11. An affine variety $V(I)$ is irreducible if and only if $I(V(I))$ is a prime ideal.
12. (Theorem of Rees) If $I = (x_1, \dots, x_n)$ is generated by a R -regular sequence and $F \in R[X_1, \dots, X_n]$ is a homogeneous polynomial of degree s such that $F(x_1, \dots, x_n) \in I^{s+1}$, then the coefficients of F are all in I . (Reduce to the case where F is a monomial.)
13. Let $I = (x_1, \dots, x_n)$ be generated by a regular sequence. Recall that the associated graded ring of I is $\text{gr}_I(R) = \bigoplus_{i=0}^{\infty} I^i/I^{i+1}$, an \mathbb{N} -graded ring with R/I in degree zero. Prove that the map $\varphi: (R/I)[X_1, \dots, X_n] \rightarrow \text{gr}_I(R)$ is an isomorphism. (It suffices to check surjectivity and injectivity on homogeneous elements; use the previous exercise for injectivity.)

14. Use the previous exercise to prove the Monomial Conjecture for Cohen-Macaulay local rings.
15. Can you state and prove generalizations of the previous three exercises that allow you to prove the Monomial Conjecture if R is only assumed to have a module M such that x_1, \dots, x_n is M -regular?
16. Let R be a local ring of dimension d and I an ideal. Prove that $\text{height } I \geq i$ if and only if I contains elements x_1, \dots, x_i forming part of a system of parameters for R .
17. Let R be Noetherian and M finitely generated. Prove that $x \in R$ is a zerodivisor on M if and only if $x \in \mathfrak{p}$ for some $\mathfrak{p} \in \text{Ass}_R M$.
18. Use the definition of the Koszul complex to write down the Koszul complex on 2 and 3 elements (in terms of matrices).
19. Let $x, y \in R$. Prove that the homology of $K_\bullet(x, y; R)$ is
 - (a) $R/(x, y)$ in degree 0;
 - (b) $\text{Ann}_R(x, y)$ in degree 2;
 - (c) $((y) :_R x)/(y)$ in degree 1 if we assume that y is a nonzerodivisor; in particular this is zero if and only if x is a nonzerodivisor on $R/(y)$.
20. Compute the intersection multiplicity of two randomly chosen plane curves at the origin, for example $y = x^3$ and $y = 0$.
21. Let $I = (x^3 - w^2y, x^2z - wy^2, xy - wz, y^3 - xz^2)$ and $J = (w, z)$ in $\mathbb{C}[x, y, z, w]$. Show that the length of the tensor product is 5, but the intersection multiplicity is 4.
22. Let $R = k[x_1, \dots, x_n]$ with k a field of characteristic p such that $[k : k^p] < \infty$. Prove that R is a finitely generated free module over the subring R^p of p^{th} powers.
23. If $R \subseteq S$ are rings such that R is a direct summand of S , prove that $IS \cap R = I$ for every ideal I of R .
24. If R is a domain which is a direct summand of every module-finite extension, prove that R is integrally closed. (Apply the previous exercise to the extension $R \subset R[\frac{y}{x}]$ where $\frac{y}{x}$ is an integral element.)
25. (For those who know about Gorenstein rings.) Let (R, \mathfrak{m}) be a Gorenstein local ring and \underline{x} a system of parameters. Let $\Delta \in R$ be a lift of a generator for the socle of $R/(\underline{x})$. Let M

be an R -module and assume \underline{x} is M -regular. Prove that R splits as a direct summand from M if and only if $\Delta M \not\subseteq \underline{x}M$. (The forward direction is easier; for the reverse, use the fact that $\varprojlim R/(x_1^t, \dots, x_d^t)$, where the maps are all multiplication by $x_1 \cdots x_d$, is isomorphic to the injective hull $E_R(k)$.)

26. Let R be a ring, I a finitely generated ideal, and M an R -module. Prove that $\text{grade}(I, M) \geq 1$ if and only if $\text{Hom}_R(R/I, M) = 0$, if and only if I annihilates some element of M . When R is Noetherian, these are equivalent to existence of some $\mathfrak{p} \in \text{Ass}_R M$ with $I \subseteq \mathfrak{p}$.